

Climate Model Report Card

Model Name: Community Earth System Model version 1.0 with Community Atmospheric Model version 5

Developers: National Science Foundation, US Department of Energy, and National Center for Atmospheric Research (NSF-DOE-NCAR)

Data Portal: <https://cera-www.dkrz.de>, www.earthsystemgrid.org

Spatial Resolution: 0.94° x 1.25°

Temporal Resolution: monthly

LAKE COMPONENT

Name: Lake model within NCAR's Community Land Model, version 4 (CLM4.0)

Reference: ¹Oleson, K. W., Dai, Y., Bonan, G., Bosilovich, M., Dickinson, R., Dirmeyer, P., ... & Zeng, X. (2010). Technical description of version 4.0 of the Community Land Model (CLM). *Tech. Note NCAR/TN-478+STR*.

Description: The lake model is from Zeng et al. (2002), which is based on various one-dimensional models that vertically solve the thermal diffusion equation for 10 layers of water and ice. Each grid cell is assigned a percentage of lake, wetland, glacier, urban, and soil, where lake percentages are from Cogley's (1991) 1.0° x 1.0° perennial freshwater lake data. Snow cover above lakes is greatly simplified and soil beneath lakes is not considered.¹ CLM4.0's lake model was greatly improved for deep lakes with CLM-LISSS (Lake, Ice, Snow, and Sediment Simulator), but this was not used in CMIP5.²

Vertical Layers | Depths: 10 layers for a maximum depth of 50m

Vertical Mixing (y/n): Yes

Horizontal Mixing (y/n): No

Lake Ice: The 10 layers can be water or ice, and frozen lake albedos are based on sea ice values from NCAR LSM (Bonan, 1996).¹

ATMOSPHERE COMPONENT

Name: Community Atmosphere Model, version 5.0 (CAM5)

Reference: ³Neale, R. B., Chen, C. C., Gettelman, A., Lauritzen, P. H., Park, S., Williamson, D. L., ... & Taylor, M. A. (2012). Description of the NCAR Community Atmosphere Model (CAM 5.0). *Tech. Note NCAR/TN-486+STR*.

Physical Parameterizations: Deep convection scheme (Zhang & McFarlane, 1995) modified for convective momentum transports (Richter & Rasch, 2008), shallow convection schemes (Park & Bretherton, 2009), stratiform cloud macrophysics (Neale et al., 2012), cloud microphysics (Morrison & Gettelman, 2008; Gettelman et al., 2008), correlated-k distribution Rapid Radiative Transfer Model (Iacono et al., 2008), statistical subgrid-scale cloud overlap (Pincus & Morcrette, 2003), moisture turbulence scheme (Bretherton & Park, 2009)^{3,4}

Chemistry: Modal aerosol treatment for black and organic carbon, dust, sea salt, and sulfate aerosols (Liu et al., 2012; Ghan et al., 2012). New aerosol parameterizations (compared to CAM4) allow for the indirect effect of aerosols to be estimated.⁴

Historical Run(s): 1850-2005

Future Scenario(s): RCP2.6, RCP4.5, RCP6.0, RCP8.5

Future Time Period(s): 2006-2300 (RCP2.6, RCP4.5), 2006-2100 (RCP8.5)

LAND COMPONENT

Name: Community Land Model (CLM), version 4.0

Reference: ¹Oleson, K. W., Dai, Y., Bonan, G., Bosilovich, M., Dickinson, R., Dirmeyer, P., ... & Zeng, X. (2010). Technical description of version 4.0 of the Community Land Model (CLM). *Tech. Note NCAR/TN-478+STR*.

Land Cover Types: 5 land cover types (soil, wetland, lake, urban, glacier) and 16 vegetation plant function types (PFTs), where each grid cell can contain multiple surface types and up to 4 PFTs.¹ Full PFT list in Oleson et al. (2010).

Soil Layers: 15 ground layers (+up to 5 for snow), where the upper 10 (0m to 3.8m) are hydrologically active soil layers and the lower 5 are bedrock layers (3.8m to 42m) for a total depth of 42m.^{1,5}

Soil Moisture: Vertical moisture between the 10 hydrologically active soil layers is governed infiltration, surface and sub-surface runoff, gradient division, gravity, canopy transpiration, and groundwater interactions¹

Runoff: Surface runoff consists of Dunne runoff (overland-flow from saturation excess) and Hortonian runoff (infiltration excess). A TOPMODEL-based runoff model called SIMTOP, described by Niu et al. (2005), is used for runoff parameterization.⁵

Sub-Grid Lakes (y/n): Yes. Grid cells can be assigned a percentage of lake, wetland, glacier, and soil, with lake percentages from Cogley's (1991) 1.0° x 1.0° perennial freshwater lake data.¹

Carbon Fluxes: CLM4 contains a prognostic carbon-nitrogen (CN) biogeochemical model (Thornton et al., 2007) that simulates carbon cycling and nitrogen limitation in vegetation, litter, and soil-organic matter, although it is turned off for this model (see CESM1-BGC, where it is turned on).^{1,4,6}

Land Use Change: CLM4 includes transient land cover and land-use change (LCLUC) with 0.5° resolute data from Hurtt et al. (2006) that describes changes in land cover from 1850-2005 between crop, pasture, primary vegetation, and secondary vegetation classes.^{1,5}

Groundwater: There's an unconfined aquifer below the 10 soil layers.⁵

Additional References

²Deng, B., Liu, S., Xiao, W., Wang, W., Jin, J., & Lee, X. (2013). Evaluation of the CLM4 lake model at a large and shallow freshwater lake. *Journal of Hydrometeorology*, 14(2), 636-649.

⁴Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., ... & Marshall, S. (2013). The community earth system model: a framework for collaborative research. *Bulletin of the American Meteorological Society*, 94(9), 1339-1360.

⁵Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., ... & Slater, A. G. (2011). Parameterization improvements and functional and structural advances in version 4 of the Community Land Model. *Journal of Advances in Modeling Earth Systems*, 3(1).

⁶Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., ... & Zhang, M. (2011). The community climate system model version 4. *Journal of climate*, 24(19), 4973-4991.